AD-A256 288



ELECTRONICALLY ACTIVE CYCLOCARBORANE-METAL-ARENE ASSEMBLIES

Final Report

Russell N. Grimes

July 31, 1992

U. S. ARMY RESEARCH OFFICE

Contract No. DAAL03-89-K-0105

University of Virginia

Approved for public release; distribution unlimited

92 9 30 029



REPORT DOCUMENTATION PAGE

Form Approved
OMB No 0704-0188

Davis Highway, Suite 1204 Arrington, VA 22222430	2. REPORT DATE		
1. AGENCY USE ONLY (Leave blank)	July 31, 1992	3. REPORT TYPE AND DATES COVERED Final 6/1/89-5/31/92	
4. TITLE AND SUBTITLE			5. FUNDING NUMBERS
Electronically Active Cycl	ocarborane-Metal-Aren	e Assemblies	
6. AUTHOR(S)	 		DAAL03-89-K-0105
Russell N. Grimes			
7. PERFORMING ORGANIZATION NAME	(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION
University of Virginia			REPORT NUMBER
Charlottesville, Virgin			
, 0			
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSORING / MONITORING
U. S. Army Research Off	ice		AGENCY REPORT NUMBER
P. O. Box 12211	07700 0011		
Research Triangle Park,	NC 27709-2211		}
11. SUPPLEMENTARY NOTES			<u> </u>
The view, opinions and	or findings contain	ned in this repo	ort are those of the
author(s) and should no			
position, policy, or de			
12a. DISTRIBUTION / AVAILABILITY STA			12b. DISTRIBUTION CODE
Approved for public re	lease; distribution	unlimited.	
<u></u>			1

13. ABSTRACT (Maximum 200 words)

The designed synthesis of stable, soluble macromolecular complexes incorporating MC_2B_3 subunits (in which M is a transition metal) was explored in detail, with emphasis on (1) the efficient preparation of viable building-block molecules such as $Cp*Co(Et_2C_2B_3H_5)$ ($Cp*=\eta^5-C_5Me_5$); (2) the derivatization and tailoring of these via attachment of substituents (including organic functional groups) to the carborane rings; (3) the construction of multidecker stacks via a novel metal-promoted stacking process; (4) the linkage of stacked sandwiches via coupling, fusion, and other methods; and (5) initial studies on the preparation of electroactive or magnetoactive solid state materials from metallacarborane precursors. The synthesis of novel metallacarborane sandwiches that are direct structural analogues of metallocenes, and mixed-sandwich salts of the type Cp_2M^+ [$(Et_2C_2B_n)_2M'$] in which M and M' are transition metals was also achieved. Detailed studies of the molecular and electronic structures of many of the above systems were conducted via NMR, ESR, X-ray crystallography, and electrochemical methods.

metallacarboranes sar	tal-arene complexes idwich complexes iltidecker sandwich comple	macromolecules organometallic polymers xes	15. NUMBER OF PAGES 16. PRICE CODE
17: SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT
UNCLASSIFIED	UNCLASSIFIED	UNCLASSIFIED	UL

NSN 7540-01-280-5500

I. Statement of Problem

This research was concerned with the development of designed synthetic routes to stable multi-metal stacked and linked sandwich complexes that are potential precursors to electroactive or magnetoactive polymers or solid-state materials, as well as studies of their electronic and molecular structures, physical properties, and chemical reactivity.

II. Summary of Important Results

1. Controlled derivitization of nido-LM(RR' $C_2B_3H_5$) synthons (LM = Cp*Co, CpCo, (arene)Fe, or (arene)Ru). Procedures we developed earlier for alkylating the C_2B_3 rings in these complexes have been extended to generate a wide variety of organic and halogen derivatives, primarily of the cobalt species $Cp*Co(Et_2C_2B_3H_5)$.

A method for selective derivatization at the cage carbon atoms via desilylation of C-trimethylsilyl closo clusters was found, that surprisingly involves catalytic cleavage of the C-Si bonds, apparently unprecedented, via the action of $(C_4H_9)_4N^+F^-$ (TBAF). The cage C-H protons in the parent complex are acidic and can be replaced by organic substituents as shown.

2. Directed Synthesis of Linked Sandwich Complexes. The above methods for attaching functional groups regionselectively on the C_2B_3 ring were exploited to prepare hydrocarbon-linked oligomers via reactions of organic dihalides:


Alternatively, we have also shown that polycyclic hydrocarbon ligands can be employed to link multisandwich species via metal- C_n ring coordination:

3. Synthesis of Tetradecker Sandwich Complexes via Metal-Promoted Stacking of Double-Decker Synthons. Although we had previously tried this approach unsuccessfully, in 1991 we found that tetradecker formation via stacking of metallacarborane double-decker anions can be achieved when the "middle" hydrogen [B(5)-H] is replaced by a suitable substituent, e.g., Cl, Br, acetyl, methyl; the parent (X = H) complex can also be isolated provided exposure to air and silica are avoided. The tetradecker sandwich geometry has been confirmed by X-ray structural studies on several Co-Co-Co, Co-Ni-Co, and Co-Ru-Co complexes.

$$X = H, Et, CH_2Ph, CH_2C_6F_5, CH_2OMe$$


O B, BH • C X = C(O)Me, CI, Br, CH₂C≡CMe M = Co, Ni, Ru

4. Designed Synthesis of Multi-Tetradecker Sandwich Oligomers and Polymers. The tetradecker stacking reaction has been employed to prepare a number of macromolecular polyarene-linked complexes such as those illustrated (the 5-cobalt species has been confirmed by X-ray crystallography). Although the phenylene-linked systems are unlikely to have delocalization of electrons between the stacks, we have recently prepared analogous systems employing fulvalene $[(C_5H_4)_2]$ as the linking unit, and the prospects for delocalization (hence conductivity) are good.



5. Studies of Electronic Structures and Properties Multinuclear NMR (¹H, ¹¹B, ¹³C), ESR, cyclic voltammetry, Mössbauer, magnetic susceptibility studies, and X-ray crystallographic structure determinations have been employed to probe the systems described above, in some cases in collaboration with other research groups. These complexes undergo multiple oxidation and reduction processes without degradation or irreversible structural change, in contrast to many hydrocarbon-metal sandwich species (most of which, other than ferrocene, tend toward oxidative instability).

The redox stability of small-carborane stabilized metal sandwich complexes is well documented, as for example, in the reversible oxidation of Fe^{II}-arenes (otherwise virtually unknown) and in the ring-migration of Fe on indenyl-ferracarborane sandwiches.



6. Coupling and Fusion of Metallacarborane Substrate Molecules. As outlined in several recent Progress Reports, a number of modes of coupling and linkage of small cobaltacarboranes (some of them novel) have been found in recent work. In some cases the reactions involved are analogous to classical procedures (e.g. Wurtz coupling of halo-substituted species), while in others the processes involve metal-promoted oxidative fusion and are similar to fusions discovered earlier in our group.

III. Publications produced under this Contract

- J. H. Davis, Jr., M. D. Attwood, and R. N. Grimes, "Organotransition-Metal Metalla-carboranes. 15. Regiospecific B-Alkylation of (arene)M($\text{Et}_2\text{C}_2\text{B}_3\text{H}_5$)(M = Fe, Ru) and (C₅Me₅)Co($\text{Et}_2\text{C}_2\text{B}_3\text{H}_5$) Sandwich Complexes," Organometallics, 1990, 9, 1171.
- M. D. Attwood, J. H. Davis, Jr., and R. N. Grimes, "Organotransition-Metal Metallacarboranes. 16. Directed Synthesis of Arene-Linked Metal Sandwich Complexes via Controlled Substitution on Boron," <u>Organometallics</u>, 1990, 9, 1177.
- J. H. Davis, Jr., M. A. Benvenuto, and R. N. Grimes, "Organotransition-Metal Metallacarboranes. 18. η⁶, η⁵-Benzyltetramethylcyclopentadieneide(1-) as a Bridging Ligand in Multilevel Iron-Cobalt Sandwich Complexes," <u>Inorg. Chem.</u> **1991**, 30, 1765.

continued

- K. W. Piepgrass, J. H. Davis, Jr., M. Sabat, and R. N. Grimes, "Unusual Organic Chemistry on a Metallacarborane Substrate: Formation of a B-Vinyl Ester from Acetyl Chloride", J. Am. Chem. Soc., 1991, 113, 680.
- K. W. Piepgrass, J. H. Davis, Jr., M. Sabat, and R. N. Grimes, "Electronic Control of Metallacarborane Stacking Reactions. Directed Synthesis of Cp*Co(C₂B₃)M(C₂B₃)CoCp* Tetradecker Sandwiches", <u>J. Am. Chem. Soc.</u>, 1991, 113, 681.
- R. N. Grimes, "Small Carboranes as Building Blocks in Designed Organometallic Synthesis," <u>Pure Appl. Chem.</u>, 1991, 63, 369.
- A. Fessenbecker, M. Stephan, R. N. Grimes, H. Pritzkow, U. Zenneck, and W. Siebert, "Organotransition-Metal Metallacarboranes. 19. Indenyliron(II) and -iron(III) Complexes and Related Species. $\eta^6 \to \eta^5$ Haptotropic Rearrangement, Electrochemistry, and Polyhedral Expansion of (Arene)Fe(Et₂C₂B₄H₄) Clusters", <u>J. Am. Chem. Soc.</u>, 1991, 113, 3061.
- M. A. Benvenuto and R. N. Grimes, "Organotransition-Metal Metallacarboranes. 20. Bu₄N*F* Fluoride-Catalyzed C-Si Bond Cleavage in Cp*Co(Me₃Si)₂C₂B₄H₄. Synthesis of Parent Cp*CoC₂B₄H₆ and Conversion to C-Substituted Cp*CoRR'C₂B₄H₄ Derivatives", Inorg. Chem. 1991, 30, 2836.
- R. N. Grimes, "Cyclocarborane-Stabilized Multidecker/Multicluster Sandwich Compounds and Linked Molecular Systems", in <u>Electron-Deficient Boron and Carbon Clusters</u>, G. A. Olah, K. Wade, and R. E. Williams, Eds., John Wiley & Sons, Inc., New York, 1991, Chapter 11, pp 261-285.
- M. Stephan, J. H. Davis, Jr., X. Meng, K. J. Chase, Jan Hauss, U. Zenneck, H. Pritzkow, W. Siebert, and R. N. Grimes, "Organotransition-Metal Metallacarboranes. 25. Redox Chemistry and Electronic Studies of Mono- and Dinuclear Iron(II)/Iron(III) Sandwich Complexes", J. Am. Chem. Soc. 1992, 114, 5214.
- K. W. Piepgrass and R. N. Grimes, "(C₅Me₅)Co(Et₂C₂B₃H₄-R) Metallacarborane Sandwich Complexes as Versatile Synthons. Introduction of Organomethyl Functional Groups at Boron", <u>Organometallics</u> 1992, <u>11</u>, 2397.
- K. W. Piepgrass, K. E. Stockman, M. Sabat, and R. N. Grimes, "(C₅Me₅)Co(Et₂C₂B₃H₄-R) Metallacarborane Sandwich Complexes as Versatile Synthons. Introduction of Electron-Withdrawing Substituents", <u>Organometallics</u> 1992, <u>11</u>, 2404.
- R. N. Grimes, "Boron-Carbon Ring Ligands in Organometallic Synthesis", Chem. Rev. 1992, 92, 251.
- M. A. Benvenuto and R. N. Grimes, "Organotransition-Metal Metallacarboranes. 28. Controlled Substitution at Cage Carbon Atoms in CoC₂B₄ Clusters. Linkage of Bis(cobaltacarborane) Sandwich Complexes via Carbon-Bound Bridging Groups", <u>Inorg. Chem.</u>, in press.
- M. A. Benvenuto, M. Sabat, and R. N. Grimes, "Organotransition-Metal Metallacarboranes. 29. Synthesis of Selectively C- and B-Substituted Double- and Triple-Decker Sandwiches. (η⁵-C₅Me₅)Co^{III}(η⁵-R₂C₂B₃R'₃H₂) Cobalticinium Analogues", <u>Inorg. Chem.</u>, in press.

K. W. Piepgrass, Xiangsheng Meng, M. Hölscher, M. Sabet and R. N. Grimes, "Tetradecker Metallacarborane Sandwiches: Synthesis via Doul 3-Decker Stacking and Structural Characterization of Co-Co-Co, Co-Ni-Co, and Co-Ru-Co Complexes", submitted for publication.

IV. Participating Scientific Personnel

Principal Investigator: Russell N. Grimes, Professor of Chemistry

Postdoctoral Associates:

Kent Piepgrass Xiangsheng Meng Simon Waterworth

Graduate Students (degrees earned):

Kevin Chase (Ph.D. August 1990) Mark Benvenuto (Ph. D. January 1992) Xiaotai Wang Kenneth Stockman